Nuclide Analysis Results of Radioactive Materials in Seawater < Coast>

Reference

(Data summarized on March 8)

Place of Sampling	North of Discharge Channel of 5-6u of 1F (approx. 30m north of 5-6u discharge channel)		Around South Discharge Channel of 1F (appox. 330m south of 1-4u Discharge Channel)		Around North Discharge Channel of 2F (Around 3,4u Discharge Channel) (approx. 10 km from 1F)		Around Iwasawa Shore of 2F (appox. 7 km south of 1,2u Discharge Channel) (appox. 16 km from 1F)		Daniela Barris I and Alexander
Time of Sampling	Mar 07, 2012 (Not sampled)		Mar 07, 2012 (Not sampled)		Mar 07, 2012 08:25 am		Mar 07, 2012 08:00 am		
Detected Nuclides (Half-life)	Density of Sample (Bq/L)	Scaling Factor (/)	Density of Sample (Bq/L)	Scaling Factor (/)	Density of Sample (Bq/L)	Scaling Factor (/)	Density of Sample (Bq/L)	Scaling Factor (/)	areas in the section 6 of the appendix 2)
I-131 (about 8 days)	-	-	-	-	ND	-	ND	-	40
Cs-134 (about 2 years)	-	-	-	-	ND	-	0.97	0.02	60
Cs-137 (about 30 years)	-	-	-	-	ND	-	ND	-	90

^{*} Density by the announcement of Reactor Regulation is stated with an amount converted from Bq/cm3 to Bq/L.

Please note that these nuclides are sometimes detected even when they are below the limits, contingent on the detector or samples.

Sampling at 2 of 4 places were canceled due to bad weather condition.


^{*} Data of other nuclides are under evaluation.

^{*} In the case that two or more kinds of nuclides exist, sum of each scaling factor to the density limit is compared with 1.


^{* &}quot;ND" means the sampled data is below measurable limit.

I-131: approx. 0.64Bq/L, Cs-134: approx. 0.86Bq/L, Cs-137: approx. 1.1Bq/L

Radioactivity Density of Seawater (Bq/L)



Radioactivity Density of Seawater at North Discharge Channel of 2F (Bq/L)



Radioactivity Density of Seawater at Iwasawa Shore 2F (Bq/L)

