Reference

Nuclides Analysis Result of the Radioactive Materials in the Seawater < Coast, Fukushima Daiichi Nuclear Power Station >

(Data summarized on November 3)

| Place of Sampling | North of Unit 5-6 Discharge Channel at Fukushima Daiichi NPS (Approx. 30m North of Unit 5-6 Discharge Channel) | | Around South Discharge Channel of Fukushima Daiichi NPS (Appox. 330m South of Unit 1-4 Discharge Channel) | | (Bq/L) (The density limit in the water outside the surrounding |
|----------------------------------|--|----------------|---|----------------|--|
| Time of Sampling | Nov 2, 2012 8:15 AM | | Nov 2, 2012 8:00 AM | | |
| Detected Nuclides (Half-life) | Density of Sample (Bq/L) | Scaling Factor | Density of Sample (Bq/L) | Scaling Factor | monitored areas is provided in section 6 of Appendix 2.) |
| I-131 (Approx. 8 days) | ND | - | ND | - | 40 |
| Cs-134 (Approx. 2 years) | ND | - | ND | - | 60 |
| Cs-137 (Approx. 30 years) | ND | - | ND | - | 90 |

^{*} The density specified by the Reactor Regulation is converted from Bq/cm3 to Bq/L.

As the detection limit may vary depending on the detectors and sample properties, there are cases where nuclides below the detection limit are detected.

^{*} Data of other nuclides is under evaluation.

^{*} In the case of more than 2 nuclides, the sum of scaling factors to density limits is compared to 1.

^{* &}quot;ND" indicates that the measurement result is below the detection limit. I-131: Approx. 0.47Bq/L, Cs-134: Approx.1.1Bq/L, Cs-137: Approx.1.4Bq/L

Reference

Nuclides Analysis Result of the Radioactive Materials in the Seawater < Coast, Fukushima Daini Nuclear Power Station >

(Data summarized on November 3)

| Place of Sampling | 2F Around the North Discharge Channel (Around Unit 3-4 Discharge Channel) (Approx. 10km from 1F) | | Around the North Side of Asamigawa (Approx. 12km South of Unit 1 & 2 Discharge Channel) (Approx. 24km from 1F) | | Density Limit Specified by the Reactor Regulation (Bq/L) (The density limit in the water outside the surrounding monitored areas is provided |
|----------------------------------|--|----------------|---|----------------|--|
| Time of Sampling | Oct 30, 2012 10:10 AM | | Oct 30, 2012 7:30 AM | | |
| Detected Nuclides (Half-life) | Density of Sample (Bq/L) | Scaling Factor | Density of Sample (Bq/L) | Scaling Factor | in section 6 of Appendix 2.) |
| I-131 (Approx. 8 days) | ND | - | ND | - | 40 |
| Cs-134 (Approx. 2 years) | ND | - | ND | - | 60 |
| Cs-137 (Approx. 30 years) | ND | - | ND | - | 90 |

^{*} The density specified by the Reactor Regulation is converted from Bq/cm³ to Bq/L.


As the detection limit may vary depending on the detectors and sample properties, there are cases where nuclides below the detection limit are detected.

^{*} Data of other nuclides is under evaluation.


^{*} In the case of more than 2 nuclides, the sum of scaling factors to density limits is compared to 1.

^{* &}quot;ND" indicates that the measurement result is below the detection limit. I-131: Approx. 0.15Bq/L, Cs-134: Approx. 0.22Bq/L, Cs-137: Approx. 0.28Bq/L


Radioactivity Density of the Seawater at the North of 1F Unit 5-6 Discharge Channel (Bq/L)



Radioactivity Density of the Seawater at 1F South Discharge Channel (Bq/L)



Radioactivity Density of the Seawater at 2F North Discharge Channel (Bq/L)



Radioactivity Density of the Seawater at Around the North of Asamigawa (Bq/L)

