Reference

Nuclides Analysis Result of the Radioactive Materials in the Seawater < Coast, Fukushima Daiichi Nuclear Power Station >

(Data summarized on December 27)

| Place of Sampling | North of Unit 5-6 Discharge Daiichi N (Approx. 30m North of Unit 5 | Channel of Fukushima NPS 1-4 Discharge Channel) | the Reactor Regulation (Bq/L) | | | | |
|----------------------------------|--|---|-------------------------------|----------------|--|--|--|
| Time of Sampling | Dec 26, 2 8:00 A | | Dec 26, 2 8:30 A | | (The density limit in the water outside the surrounding monitored areas is provided in section 6 of Appendix 2.) | | |
| Detected Nuclides (Half-life) | Density of Sample (Bq/L) | Scaling Factor | Density of Sample (Bq/L) | Scaling Factor | | | |
| I-131 (Approx. 8 days) | ND | - | ND | - | 40 | | |
| Cs-134 (Approx. 2 years) | ND | - | ND | - | 60 | | |
| Cs-137 (Approx. 30 years) | ND | - | ND | - | 90 | | |

^{*} The density specified by the Reactor Regulation is converted from Bq/cm³ to Bq/L.

I-131: Approx. 0.44Bq/L, Cs-134: Approx. 1.1Bq/L, Cs-137: Approx. 1.4Bq/L

As the detection limit may vary depending on the detectors and sample properties, there are cases where nuclides below the detection limit are detected.

^{*} Data of other nuclides is under evaluation.

^{*} In the case of more than 2 nuclides, the sum of scaling factors to density limits is compared to 1.

 $[\]ensuremath{^{*}}$ "ND" indicates that the measurement result is below the detection limit.

Reference

Nuclides Analysis Result of the Radioactive Materials in the Seawater < Coast, Fukushima Daiichi Nuclear Power Station, Remeasurement >

| Place of Sampling | North of Unit 5-6 Discharge Daiichi N (Approx. 30m North of Unit 5 | IPS | Around South Discharge C Daiichi N (Appox. 330m South of Unit | IPS | the Reactor Regulation (Bq/L) | | |
|----------------------------------|--|----------------|---|--|-------------------------------|--|--|
| Time of Sampling | Nov 19, 2 8:30 A | | Nov 19, 2 8:00 A | (The density limit in the water outside the surrounding monitored areas is provided in | | | |
| Detected Nuclides (Half-life) | Density of Sample (Bq/L) | Scaling Factor | Density of Sample (Bq/L) | Scaling Factor | section 6 of Appendix 2.) | | |
| Cs-134 (Approx. 2 years) | 0.41 | 0.01 | 0.37 | 0.01 | 60 | | |
| Cs-137 (Approx. 30 years) | 0.69 | 0.01 | 0.62 | 0.01 | 90 | | |

^{*} The density specified by the Reactor Regulation is converted from Bq/cm³ to Bq/L.

^{*} In the case of more than 2 nuclides, the sum of scaling factors to density limits is compared to 1.

^{*} Analysis results by detail analysis (Phosphomolybdic acid ammonium adsorption sampling method) are noted.

^{*} Analyzed by : Tokyo Electric Power Environmental Engineering Co., Inc.

Reference

Nuclides Analysis Result of the Radioactive Materials in the Seawater < Coast, Fukushima Daini Nuclear Power Station >

(Data summarized on December 27)

| Place of Sampling | 2F Around the North D (Around Unit 3-4 Disc (Approx. 10km | charge Channel) | Around the North Sid (Approx. 12km South of U Channo (Approx. 24km | Density Limit Specified by the Reactor Regulation (Bq/L) | | | |
|----------------------------------|---|-----------------|---|--|--|--|--|
| Time of Sampling | Nov 20, 2 10:00 A | | Nov 20, 2 7:30 A | | (The density limit in the water outside the surrounding monitored areas is provided in | | |
| Detected Nuclides (Half-life) | Density of Sample (Bq/L) | Scaling Factor | Density of Sample (Bq/L) | Scaling Factor | section 6 of Appendix 2.) | | |
| I-131 (Approx. 8 days) | ND | - | ND | - | 40 | | |
| Cs-134 (Approx. 2 years) | 0.19 | 0.00 | 0.13 | 0.00 | 60 | | |
| Cs-137 (Approx. 30 years) | 0.30 | 0.00 | 0.22 | 0.00 | 90 | | |

^{*} The density specified by the Reactor Regulation is converted from Bq/cm³ to Bq/L.

I-131: Approx. 0.45Bq/L

As the detection limit may vary depending on the detectors and sample properties, there are cases where nuclides below the detection limit are detected.

^{*} Data of other nuclides is under evaluation.

^{*} In the case of more than 2 nuclides, the sum of scaling factors to density limits is compared to 1.

^{* &}quot;ND" indicates that the measurement result is below the detection limit.

^{*} As to Cs-134 and Cs-137, analysis results by detail analysis (Phosphomolybdic acid ammonium adsorption sampling method) are noted. Analyzed by Tokyo Electric Power Environmental Engineering Co., Inc.

Nuclides Analysis Result of the Radioactive Materials in the Seawater < Offshore 1/3 >

| Place of Sampling (Place No.) | 3km Off | shore of C | daka Ward (T-1 | *1 14) | 3km Off | lkedo River (T-[| 3km Offshore | of Fukush | *2 S (T-D5) | Density Limit Specified by the Reactor Regulation | | | |
|----------------------------------|--------------------------------|----------------------------|--------------------------------|----------------------------|--------------------------------|----------------------------|--------------------------------|----------------------------|--------------------------------|---|--------------------------------|-------------------|---|
| | Upper La | Upper Layer Lower Layer | | | | ayer | Lower La | ayer | Upper La | ayer | Lower La | ayer | (Bq/L) |
| Time of Sampling | Nov 13, 2 9:05 Al | | , | Nov 13, 2012 Nov | | Nov 19, 2012 8:50 AM | | Nov 19, 2012 8:50 AM | | 2012 M | Nov 19, 2012 8:20 AM | | (The density limit in the water outside the surrounding monitored |
| Detected Nuclides (Half-life) | Density of Sample (Bq/L) | Scaling Factor (/) | Density of Sample (Bq/L) | Scaling Factor | Density of Sample (Bq/L) | Scaling Factor | areas is provided in section 6 of Appendix 2.) |
| Cs-134 (Approx. 2 years) | 0.030 | 0.00 | 0.031 | 0.00 | 0.035 | 0.00 | 0.025 | 0.00 | 0.046 | 0.00 | 0.044 | 0.00 | 60 |
| Cs-137 (Approx. 30 years) | 0.050 | 0.00 | 0.051 | 0.00 | 0.052 | 0.00 | 0.042 | 0.00 | 0.080 | 0.00 | 0.080 | 0.00 | 90 |

| Place of Sampling (Place No.) | 3km Offshore | e of Fukusl | nima Daini NPS | *2 5 (T-D9) | 15km Offshor | shima Daiichi N | 3km Offs | hore of Iwa | *1 -11) | Density Limit Specified by the Reactor Regulation | | | |
|----------------------------------|--------------------------------|----------------------------|--------------------------------|----------------------------|--------------------------------|----------------------------|--------------------------------|-------------------|--------------------------------|---|--------------------------------|-------------------|---|
| | Upper La | ayer | Lower La | ayer | Upper La | ayer | Lower La | ayer | Upper La | ayer | Lower La | ayer | (Bq/L) |
| Time of Sampling | Nov 21, 2 9:10 A | - | Nov 21, 2 9:10 Al | - | Nov 15, 2012 9:00 AM | | Nov 15, 2012 9:00 AM | | Nov 15, 2012 10:35 AM | | Nov 15, 2012 10:35 AM | | (The density limit in the water outside the surrounding monitored |
| Detected Nuclides (Half-life) | Density of Sample (Bq/L) | Scaling Factor (/) | Density of Sample (Bq/L) | Scaling Factor (/) | Density of Sample (Bq/L) | Scaling Factor (/) | Density of Sample (Bq/L) | Scaling Factor | Density of Sample (Bq/L) | Scaling Factor | Density of Sample (Bq/L) | Scaling Factor | areas is provided in section 6 of Appendix 2.) |
| Cs-134 (Approx. 2 years) | 0.033 | 0.00 | 0.031 | 0.00 | 0.0047 | 0.00 | 0.0056 | 0.00 | 0.096 | 0.00 | 0.072 | 0.00 | 60 |
| Cs-137 (Approx. 30 years) | 0.061 | 0.00 | 0.056 | 0.00 | 0.0090 | 0.00 | 0.0093 | 0.00 | 0.16 | 0.00 | 0.11 | 0.00 | 90 |

^{*} The density specified by the Reactor Regulation is converted from Bq/cm³ to Bq/L.

^{*} In the case of more than 2 nuclides, the sum of scaling factors to density limits is compared to 1.

^{*} Analysis results by detail analysis (Phosphomolybdic acid ammonium adsorption sampling method) are noted.

^{*} Analyzed by: *1 THE GENERAL ENVIRONMENTAL TECHNOS Co., LTD., *2 Tokyo Electric Power Environmental Engineering Co., Inc.

Nuclides Analysis Result of the Radioactive Materials in the Seawater < Offshore 2/3 >

| Place of Sampling (Place No.) | | 1km Offshore of Nida River (T-13-1) | | | | | of Soma (T-22) | | | offshore of | <u>'</u> | Density Limit Specified by the Reactor Regulation | |
|----------------------------------|--------------------------------|-------------------------------------|--------------------------------|----------------------------|--------------------------------|-------------------|--------------------------------|-------------------|--------------------------------|-------------------|--------------------------------|---|---|
| | Upper La | ayer | Lower La | ayer | Upper La | ayer | Lower La | ayer | Upper La | ayer | Lower La | ayer | (Bq/L) |
| Time of Sampling | Nov 22, 2 6:40 A | | Nov 22, 2012 6:40 AM | | Nov 22, 2012 7:50 AM | | Nov 22, 2012 7:50 AM | | Nov 22, 2012 7:20 AM | | Nov 22, 2012 7:20 AM | | (The density limit in the water outside the surrounding monitored |
| Detected Nuclides (Half-life) | Density of Sample (Bq/L) | Scaling Factor | Density of Sample (Bq/L) | Scaling Factor (/) | Density of Sample (Bq/L) | Scaling Factor | areas is provided in section 6 of Appendix 2.) |
| Cs-134 (Approx. 2 years) | 0.028 | 0.00 | 0.024 | 0.00 | 0.017 | 0.00 | 0.013 | 0.00 | 0.027 | 0.00 | 0.017 | 0.00 | 60 |
| Cs-137 (Approx. 30 years) | 0.046 | 0.00 | 0.042 | 0.00 | 0.030 | 0.00 | 0.025 | 0.00 | 0.045 | 0.00 | 0.033 | 0.00 | 90 |

| Place of Sampling (Place No.) | Around 3km Offshore of Ukedo River (T-S3) Upper Layer Lower Layer | | | | Around 3km C | (T- | Fukushima Da S4) Lower La | | Around 2k | | Density Limit Specified by the Reactor Regulation (Bq/L) | | |
|----------------------------------|--|----------------------------|--------------------------------|----------------------------|--------------------------------|-------------------------|---------------------------------|-------------------------|--------------------------------|-------------------|--|-------------------|---|
| Time of Sampling | Nov 20, 2 6:30 A | - | / | Nov 20, 2012 6:30 AM | | Nov 20, 2012 7:00 AM | | Nov 20, 2012 7:00 AM | | 2012 M | Nov 14, 2012 1:25 PM | | (The density limit in the water outside the surrounding monitored |
| Detected Nuclides (Half-life) | Density of Sample (Bq/L) | Scaling Factor (/) | Density of Sample (Bq/L) | Scaling Factor (/) | Density of Sample (Bq/L) | Scaling Factor | Density of Sample (Bq/L) | Scaling Factor | Density of Sample (Bq/L) | Scaling Factor | Density of Sample (Bq/L) | Scaling Factor | areas is provided in section 6 of Appendix 2.) |
| Cs-134 (Approx. 2 years) | 0.032 | 0.00 | 0.027 | 0.00 | 0.026 | 0.00 | 0.019 | 0.00 | 0.11 | 0.00 | 0.099 | 0.00 | 60 |
| Cs-137 (Approx. 30 years) | 0.051 | 0.00 | 0.043 | 0.00 | 0.041 | 0.00 | 0.035 | 0.00 | 0.18 | 0.00 | 0.16 | 0.00 | 90 |

^{*} The density specified by the Reactor Regulation is converted from Bq/cm3 to Bq/L.

^{*} In the case of more than 2 nuclides, the sum of scaling factors to density limits is compared to 1.

^{*} Analysis results by detail analysis (Phosphomolybdic acid ammonium adsorption sampling method) are noted.

^{*} Analyzed by: THE GENERAL ENVIRONMENTAL TECHNOS Co., LTD.

Nuclides Analysis Result of the Radioactive Materials in the Seawater < Offshore 3/3 >

| Place of Sampling (Place No.) | Around 2km Of | fshore of I S | | ni NPS (T- | Around 4km | Offshore | of Kumagawa | (T-S8) | Arounmd 15k | m Offshore | Density Limit Specified by the Reactor Regulation | | |
|----------------------------------|--------------------------------|----------------------------|--------------------------------|----------------------------|--------------------------------|----------------------------|--------------------------------|----------------------------|--------------------------------|-------------------|---|-------------------|---|
| | Upper La | ayer | Lower La | ayer | Upper La | ayer | Lower La | ayer | Upper La | ayer | Lower La | ayer | (Bq/L) |
| Time of Sampling | Nov 14, 2 12:50 P | | Nov 14, 2 12:55 P | | Nov 18, 2 6:40 A | | Nov 18, 2012 6:45 AM | | Nov 16, 2012 7:25 AM | | Nov 16, 2012 7:30 AM | | (The density limit in the water outside the surrounding monitored |
| Detected Nuclides (Half-life) | Density of Sample (Bq/L) | Scaling Factor (/) | Density of Sample (Bq/L) | Scaling Factor | Density of Sample (Bq/L) | Scaling Factor | areas is provided in section 6 of Appendix 2.) |
| Cs-134 (Approx. 2 years) | 0.13 | 0.00 | 0.14 | 0.00 | 0.058 | 0.00 | 0.052 | 0.00 | 0.0049 | 0.00 | 0.0045 | 0.00 | 60 |
| Cs-137 (Approx. 30 years) | 0.21 | 0.00 | 0.23 | 0.00 | 0.094 | 0.00 | 0.085 | 0.00 | 0.0088 | 0.00 | 0.0099 | 0.00 | 90 |

| Place of Sampling (Place No.) | | Around 18km Offshore of Ukedo River (T-B2) Upper Layer Lower Layer | | | | Upper Layer Lower Layer | | | | | Upper Layer Lower Layer | | | | |
|----------------------------------|--------------------------------|---|--------------------------------|----------------------------|--------------------------------|-------------------------|--------------------------------|-------------------|--------------------------------|-------------------|--------------------------------|----------------------------|--|--|--|
| Time of Sampling | Nov 16, 2 6:25 A | 2012 | Nov 16, 2 6:35 A | .012 | | Оррег Сауег | | | | | | | (Bq/L) (The density limit in the water outside the surrounding monitored | | |
| Detected Nuclides (Half-life) | Density of Sample (Bq/L) | Scaling Factor | Density of Sample (Bq/L) | Scaling Factor (/) | Density of Sample (Bq/L) | Scaling Factor | Density of Sample (Bq/L) | Scaling Factor | Density of Sample (Bq/L) | Scaling Factor | Density of Sample (Bq/L) | Scaling Factor (/) | areas is provided in section 6 of Appendix 2.) | | |
| Cs-134 (Approx. 2 years) | 0.0078 | 0.00 | 0.028 | 0.00 | | | | | | | | | 60 | | |
| Cs-137 (Approx. 30 years) | 0.013 | 0.00 | 0.046 | 0.00 | | | | | | | | | 90 | | |


^{*} The density specified by the Reactor Regulation is converted from Bq/cm3 to Bq/L.

^{*} In the case of more than 2 nuclides, the sum of scaling factors to density limits is compared to 1.


^{*} Analysis results by detail analysis (Phosphomolybdic acid ammonium adsorption sampling method) are noted.

^{*} Analyzed by: THE GENERAL ENVIRONMENTAL TECHNOS Co., LTD.

Radioactivity Density of the Seawater at 1F Units 5-6 North Discharge Channel (Bq/L)




Radioactivity Density of the Seawater at 1F South Discharge Channel (Bq/L)



Sampling was conducted at around South Discharge Channel of Fukushima Daiichi NPS (appox. 330m south of Units 1-4 Discharge Channel) until November 25, 2012.

Radioactivity Density of the Seawater at 2F North Discharge Channel (Bq/L)



Radioactivity Density of the Seawater at Around the North of Asamigawa (Bq/L)

