Reference

Nuclides Analysis Result of the Radioactive Materials in the Air at the Upper Part of Unit 2 Reactor Building

(Data summarized on March 15)

Place of Sampling	Upper Part of Unit 2 Reactor Building (The Center of the Blow-out Panel, West Side Upper)		Upper Part of Unit 2 Reactor Building (The Center of the Blow-out Panel, West Side Lower)		Upper Part of Unit 2 Reactor Building (The Center of the Blow-out Panel, West Side Upper)		Density Limit Specified by the Reactor Regulation (Bq/cm³) (Density limit in the air which radiation workers breathe in
Time of Sampling	Mar 6, 2013 8:35 AM - 10:35 AM		Mar 6, 2013 8:35 AM - 10:35 AM		Mar 6, 2013 8:35 AM - 10:35 AM		
Detected Nuclides (Half-life)	Density of Sample (Bq/cm ³)	Scaling Factor (/)	Density of Sample (Bq/cm ³)	Scaling Factor (/)	Density of Sample (Bq/cm ³)	Scaling Factor (/)	is specified in section 4 of Appendix 2)
I-131 (Approx. 8 days)	ND	-	ND	-	ND	-	1E-03
Cs-134 (Approx. 2 years)	7.7E-06	0.00	ND	-	ND	1	2E-03
Cs-137 (Approx. 30 years)	9.8E-06	0.00	ND	-	ND	-	3E-03

^{*} The radioactivity density is the sum of the volatile nuclides density and the particulate nuclides density.

O.OE - O is the same as O.O x 10^{-O}

Data of other nuclides is under examination.

The detection limits are as follows. Volatile: I-131: Approx. 2E-6Bq/cm3, Cs-134: Approx.5E-6Bq/cm3, Cs-137: Approx.6E-6Bq/cm3
Particulate: I-131: Approx. 1E-6Bq/cm3, Cs-134: Approx.3E-6Bq/cm3, Cs-137: Approx.4E-6Bq/cm3
As the detection limit may vary depending on the detectors and sample properties, there are cases where nuclides below the detection limit are detected.

^{*} In the case of more than 2 nuclides, the sum of scaling factors to density limits is compared to 1.

^{* &}quot;ND" indicates that the measurement result is below the detection limit.