Underground Reservoir Nuclide Analysis Results (As of October 8, 2014)

| | | Underground Reservoir (Drain hole water) | | | | | | | | | | | | | |
|------------------------------|--|--|-----------|---------|---------|---------|---------|---------|-----------|---------|---------|---------|---------|-------------|-----------|
| | i | | ii | | iii | | iv | | V | | vi | | \ | v ii | |
| | | | Southwest | | | | | | Southwest | | | | | | Southwest |
| | | side | side | side | side | side | side | side | side | side | side | side | side | side | side |
| Sampled time | | 7:50 AM | 7:56 AM | 8:17 AM | 8:03 AM | 8:25 AM | 8:12 AM | 9:13 AM | 9:18 AM | 8:52 AM | 8:48 AM | 9:04 AM | 8:55 AM | 8:44 AM | 8:40 AM |
| Chloride concentration (ppm) | | 9 | 8 | 1 | 6 | 2 | 2 | 11 | 5 | 9 | 7 | 10 | 2 | 5 | 9 |
| Radioactive concentration | I-131 | <2.5E-2 | <2.4E-2 | <2.7E-2 | <2.6E-2 | <2.1E-2 | <2.1E-2 | <2.7E-2 | <2.1E-2 | <2.4E-2 | <2.4E-2 | <2.6E-2 | <2.5E-2 | <2.1E-2 | <2.4E-2 |
| | Cs-134 | <5.1E-2 | <3.5E-2 | <4.2E-2 | <3.9E-2 | <4.1E-2 | <3.7E-2 | <4.0E-2 | <4.2E-2 | <4.1E-2 | <3.9E-2 | <4.5E-2 | <4.0E-2 | <4.6E-2 | <4.3E-2 |
| | Cs-137 | <6.3E-2 | <5.6E-2 | <6.4E-2 | <5.7E-2 | <6.3E-2 | <6.0E-2 | <6.5E-2 | <5.7E-2 | <6.5E-2 | <5.7E-2 | <6.5E-2 | <5.6E-2 | <6.3E-2 | <5.8E-2 |
| | γ nuclides other than the major 3 nuclides | ND | ND | ND | ND | ND | ND | ND | ND | ND | ND | ND | ND | ND | ND |
| (Bq/cm ³) | ΑΙΙ β | 2.8E-1 | <3.0E-2 | <3.0E-2 | <3.0E-2 | <3.0E-2 | <3.0E-2 | <3.0E-2 | <3.0E-2 | <3.0E-2 | 7.1E-2 | <3.0E-2 | <3.0E-2 | <3.0E-2 | <3.0E-2 |

Half-life period I-131: Approx. 8 days, Cs-134: Approx. 2 years, Cs-137: Approx. 30 years

| | | Underground Reservoir (Leakage detector hole water) | | | | | | | | | | | | | | |
|------------------------------|--|---|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--|
| | | i | | ii | | iii | | iv | | V | | vi | | vii / | | |
| | | Northeast side | Southwest side | Northeast side | Southwest side | Northeast side | Southwest side | Northeast side | Southwest side | Northeast side | Southwest side | Northeast side | Southwest side | Northeast side | Southwest side | |
| Sampled time | | 7:47 | 8:00 AM | 7:43 AM | 8:07 AM | 8:21 AM | 8:30 AM | 9:15 AM | Not sampled | / | / | 8:59 AM | Not sampled | | | |
| Chloride concentration (ppm) | | 10 | 6 | 10 | 10 | 7 | 7 | 12 | | | | 5 | | | | |
| Radioactive concentration | I-131 | <2.4E-2 | <2.4E-2 | <2.8E-2 | <2.8E-2 | <2.2E-2 | <2.1E-2 | <2.2E-2 | | | | <2.7E-2 | | / | | |
| | Cs-134 | <4.4E-2 | <4.1E-2 | <4.1E-2 | <5.6E-2 | <4.2E-2 | <3.8E-2 | <4.4E-2 | | | | <3.9E-2 | | | | |
| | Cs-137 | <6.8E-2 | <5.7E-2 | <6.6E-2 | <6.0E-2 | <6.3E-2 | <5.8E-2 | <6.4E-2 | | | | <6.4E-2 | | | | |
| | γ nuclides other than the major 3 nuclides | ND | ND | ND | ND | ND | ND | ND | | | | ND | | | | |
| (Bq/cm ³) | ΑΙΙ β | 6.6E+1 | 3.5E-2 | 1.6E+1 | <3.0E-2 | 1.5E+1 | 5.0E+0 | <3.0E-2 | | | | <3.0E-2 | | | | |

Half-life period I-131: Approx. 8 days, Cs-134: Approx. 2 years, Cs-137: Approx. 30 years

(Note 1) O.OE±O is the same as O.O x 10^{±O}.

(Note 2) The figures written next to "<" indicate the detection limit when the measurement result is below the detection limit.

(Note 3) "ND" indicates that the measurement result of y nuclides other than the major 3 nuclides are below the detection limit.