Nuclide Analysis Results of Fish < Port Area of Fukushima Daiichi Nuclear Power Station> < 1/4>

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)			
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)	
Greenling (muscle) No.1	Inner port (Near shallow draft quay) in Fukushima Daiichi NPS	Dec 23, 2014	330	1,100	1,430	
Sea raven (muscle) No.1	Inner port (Near east breakwater) in Fukushima Daiichi NPS	Dec 11, 2014	20	80	100	
Flatfish (muscle) No.1	Inner port (Near south breakwater) in Fukushima Daiichi NPS	Dec 2, 2014	13	60	73	
Brown hakeling (muscle) No.1	Inner port (Near south breakwater) in Fukushima Daiichi NPS	Dec 11, 2014	200	690	890	
Greenling (muscle) No.2	Inner port (Near port entrance) in Fukushima Daiichi NPS	Dec 9, 2014	520	1,600	2,120	
Sea raven (muscle) No.2	Inner port (Near port entrance) in Fukushima Daiichi NPS	Dec 9, 2014	22	97	119	
Marbled sole (muscle) No.1	Inner port (Near port entrance) in Fukushima Daiichi NPS	Dec 9, 2014	780	2,600	3,380	
Acanthopagrus schlegeli (muscle)	Inner port (Near port entrance) in Fukushima Daiichi NPS	Dec 12, 2014	ND (6.4)	31	31	
Flatfish (muscle) No.2	Inner port (Near port entrance) in Fukushima Daiichi NPS	Dec 12, 2014	81	230	311	
Marbled sole (muscle) No.2	Inner port (Near port entrance) in Fukushima Daiichi NPS	Dec 12, 2014	ND (6.1)	15	15	

^{*} When analyzed result is less than a detection limit level of the radioactivity concentration, showed as "ND"; as of individual Nuclide, detection limit level is showed in parenthesis.

^{*} The sum of Cs-134 and Cs-137 radioactivity concentration as a standard value (since April 1, 2012) is 100 Bq/kg.

Nuclide Analysis Results of Fish < Port Area of Fukushima Daiichi Nuclear Power Station> < 2/4>

1		T	1	·		
Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)			
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)	
Schlegel's black rockfish (muscle)	Inner port (Near port entrance) in Fukushima Daiichi NPS	Dec 15, 2014	50	150	200	
Sea raven (muscle) No.3	Inner port (Near port entrance) in Fukushima Daiichi NPS	Dec 15, 2014	ND (7.6)	54	54	
Marbled sole (muscle) No.3	Inner port (Near port entrance) in Fukushima Daiichi NPS	Dec 15, 2014	45	140	185	
Sea raven (muscle) No.4	Inner port (Near port entrance) in Fukushima Daiichi NPS	Dec 18, 2014	16	53	69	
Jacopever (muscle) No.1	Inner port (Near port entrance) in Fukushima Daiichi NPS	Dec 18, 2014	53,000	170,000	223,000	
Flatfish (muscle) No.3	Inner port (Near port entrance) in Fukushima Daiichi NPS	Dec 18, 2014	39	160	199	
Marbled sole (muscle) No.4	Inner port (Near port entrance) in Fukushima Daiichi NPS	Dec 18, 2014	28	120	148	
Sea raven (muscle) No.5	Inner port (Near port entrance) in Fukushima Daiichi NPS	Dec 20, 2014	ND (7.8)	24	24	
Common skete (muscle)	Inner port (Near port entrance) in Fukushima Daiichi NPS	Dec 20, 2014	130	380	510	
Sebastes cheni (muscle) No.1	Inner port (Near port entrance) in Fukushima Daiichi NPS	Dec 20, 2014	1,500	4,800	6,300	

^{*} When analyzed result is less than a detection limit level of the radioactivity concentration, showed as "ND"; as of individual Nuclide, detection limit level is showed in parenthesis.

^{*} The sum of Cs-134 and Cs-137 radioactivity concentration as a standard value (since April 1, 2012) is 100 Bq/kg.

Nuclide Analysis Results of Fish < Port Area of Fukushima Daiichi Nuclear Power Station > < 3/4 >

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)			
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)	
Jacopever (muscle) No.2	Inner port (Near port entrance) in Fukushima Daiichi NPS	Dec 20, 2014	9,900	33,000	42,900	
Spotbelly rockfish (muscle)	Inner port (Near port entrance) in Fukushima Daiichi NPS	Dec 20, 2014	820	2,600	3,420	
Sea raven (muscle) No.6	Inner port (Near port entrance) in Fukushima Daiichi NPS	Dec 23, 2014	46	180	226	
Sea raven (muscle) No.7	Inner port (Near port entrance) in Fukushima Daiichi NPS	Dec 24, 2014	ND (7.2)	23	23	
Alaska pollack (muscle) No.1	Inner port (Near port entrance) in Fukushima Daiichi NPS	Dec 24, 2014	ND (6.2)	ND (6.0)	ND	
Greenling (muscle) No.3	Inner port (Near port entrance) in Fukushima Daiichi NPS	Dec 25, 2014	100	310	410	
Sebastes cheni (muscle) No.2	Inner port (Near port entrance) in Fukushima Daiichi NPS	Dec 25, 2014	6,300	20,000	26,300	
Greenling (muscle) No.4	Inner port (Near port entrance) in Fukushima Daiichi NPS	Dec 26, 2014	70	220	290	
Alaska pollack (muscle) No.2	Inner port (Near port entrance) in Fukushima Daiichi NPS	Dec 26, 2014	ND (7.8)	8.8	8.8	
Alaska pollack (muscle) No.3	Inner port (Near port entrance) in Fukushima Daiichi NPS	Dec 28, 2014	ND (6.6)	ND (6.7)	ND	

^{*} When analyzed result is less than a detection limit level of the radioactivity concentration, showed as "ND"; as of individual Nuclide, detection limit level is showed in parenthesis.

^{*} The sum of Cs-134 and Cs-137 radioactivity concentration as a standard value (since April 1, 2012) is 100 Bq/kg.

Nuclide Analysis Results of Fish < Port Area of Fukushima Daiichi Nuclear Power Station > < 4/4 >

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)			
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)	
Marbled sole (muscle) No.5	Inner port (Near port entrance) in Fukushima Daiichi NPS	Dec 28, 2014	240	800	1,040	
Brown hakeling (muscle) No.2	Inner port (Near port center) in Fukushima Daiichi NPS	Dec 16, 2014	300	1,000	1,300	

^{*} When analyzed result is less than a detection limit level of the radioactivity concentration, showed as "ND"; as of individual Nuclide, detection limit level is showed in parenthesis.

^{*} The sum of Cs-134 and Cs-137 radioactivity concentration as a standard value (since April 1, 2012) is 100 Bq/kg.