

Nuclide Analysis Results of Fish and Shelfish (Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <1/11>
(excluding inside the port)

(Data summarized on January 20)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Greenling (muscle)	Around 1km Offshore of Ota River (T-S1)	Dec 20, 2014	ND (3.2)	3.8	3.8
Schlegel's black rockfish (muscle)	Around 1km Offshore of Ota River (T-S1)	Dec 20, 2014	6.9	24	30.9
Sea raven (muscle)	Around 1km Offshore of Ota River (T-S1)	Dec 20, 2014	ND (3.2)	5.0	5.0
Common skete (muscle)	Around 1km Offshore of Ota River (T-S1)	Dec 20, 2014	11	24	35
Ovalipes punctatus (whole)	Around 1km Offshore of Ota River (T-S1)	Dec 20, 2014	ND (3.5)	ND (4.3)	ND
Flatfish (muscle)	Around 1km Offshore of Ota River (T-S1)	Dec 20, 2014	ND (3.6)	ND (4.0)	ND
Greenling (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	Dec 20, 2014	ND (3.6)	3.7	3.7
Blue crab (whole)	Around 3km Offshore of Odaka Ward (T-S2)	Dec 20, 2014	ND (4.1)	ND (4.4)	ND
Snailfish (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	Dec 20, 2014	ND (3.3)	ND (3.8)	ND
Sea raven (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	Dec 20, 2014	ND (3.7)	ND (4.1)	ND

* When analyzed result is less than a detection limit level of the radioactivity concentration, showed as "ND"; as of individual Nuclide, detection limit level is showed in parenthesis.

* The sum of Cs-134 and Cs-137 radioactivity concentration as a standard value (since April 1, 2012) is 100 Bq/kg.

* Analyzed by Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shelfish (Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <2/11>
(excluding inside the port)

(Data summarized on January 20)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Common skete (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	Dec 20, 2014	4.6	14	18.6
Sebastes cheni (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	Dec 20, 2014	12	37	49
Ovalipes punctatus (whole)	Around 3km Offshore of Odaka Ward (T-S2)	Dec 20, 2014	ND (3.8)	ND (4.2)	ND
Flatfish (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	Dec 20, 2014	ND (3.2)	ND (3.9)	ND
Marbled sole (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	Dec 20, 2014	7.3	21	28.3
Stingray (muscle)	Around 3km Offshore of Ukedo River (T-S3)	Dec 5, 2014	ND (3.6)	6.9	6.9
Blue crab (whole)	Around 3km Offshore of Ukedo River (T-S3)	Dec 5, 2014	ND (3.6)	ND (4.1)	ND
Common skete (muscle)	Around 3km Offshore of Ukedo River (T-S3)	Dec 5, 2014	7.7	18	25.7
Ovalipes punctatus (whole)	Around 3km Offshore of Ukedo River (T-S3)	Dec 5, 2014	ND (3.8)	ND (4.1)	ND
Flatfish (muscle)	Around 3km Offshore of Ukedo River (T-S3)	Dec 5, 2014	ND (2.9)	ND (3.4)	ND

* When analyzed result is less than a detection limit level of the radioactivity concentration, showed as "ND"; as of individual Nuclide, detection limit level is showed in parenthesis.

* The sum of Cs-134 and Cs-137 radioactivity concentration as a standard value (since April 1, 2012) is 100 Bq/kg.

* Analyzed by Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shelfish (Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <3/11>
(excluding inside the port)

(Data summarized on January 20)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Sea robin (muscle)	Around 3km Offshore of Ukedo River (T-S3)	Dec 5, 2014	ND (3.6)	3.9	3.9
Marbled sole (muscle)	Around 3km Offshore of Ukedo River (T-S3)	Dec 5, 2014	9.6	36	45.6
Stingray (muscle)	Around 3km Offshore of Fukushima Daiichi NPS (T-S4)	Dec 5, 2014	ND (3.7)	7.1	7.1
Blue crab (whole)	Around 3km Offshore of Fukushima Daiichi NPS (T-S4)	Dec 5, 2014	ND (3.4)	ND (4.3)	ND
Angel shark (muscle)	Around 3km Offshore of Fukushima Daiichi NPS (T-S4)	Dec 5, 2014	49	160	209
Lophius litilon (whole)	Around 3km Offshore of Fukushima Daiichi NPS (T-S4)	Dec 5, 2014	ND (3.3)	ND (3.7)	ND
Common skete (muscle)	Around 3km Offshore of Fukushima Daiichi NPS (T-S4)	Dec 5, 2014	14	39	53
Microstomus achne (muscle)	Around 3km Offshore of Fukushima Daiichi NPS (T-S4)	Dec 5, 2014	15	57	72
Flatfish (muscle)	Around 3km Offshore of Fukushima Daiichi NPS (T-S4)	Dec 5, 2014	ND (4.0)	ND (3.2)	ND
Sea robin (muscle)	Around 3km Offshore of Fukushima Daiichi NPS (T-S4)	Dec 5, 2014	ND (3.6)	5.0	5.0

* When analyzed result is less than a detection limit level of the radioactivity concentration, showed as "ND"; as of individual Nuclide, detection limit level is showed in parenthesis.

* The sum of Cs-134 and Cs-137 radioactivity concentration as a standard value (since April 1, 2012) is 100 Bq/kg.

* Analyzed by Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shelfish (Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <4/11>
(excluding inside the port)

(Data summarized on January 20)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Marbled sole (muscle)	Around 3km Offshore of Fukushima Daiichi NPS (T-S4)	Dec 5, 2014	4.1	10	14.1
Lophius litilon (whole)	Around 2km Offshore of Kido River (T-S5)	Dec 12, 2014	ND (4.5)	ND (4.1)	ND
Common skete (muscle)	Around 2km Offshore of Kido River (T-S5)	Dec 12, 2014	ND (4.9)	15	15
Microstomus achne (muscle)	Around 2km Offshore of Kido River (T-S5)	Dec 12, 2014	13	33	46
Flatfish (muscle)	Around 2km Offshore of Kido River (T-S5)	Dec 12, 2014	ND (4.9)	16	16
Sea raven (muscle)	Offshore 2km vicinity of Fukushima Daiichi NPS (T-S7)	Dec 12, 2014	ND (3.8)	6.6	6.6
Common skete (muscle)	Offshore 2km vicinity of Fukushima Daiichi NPS (T-S7)	Dec 12, 2014	8.1	26	34.1
Flatfish (muscle)	Offshore 2km vicinity of Fukushima Daiichi NPS (T-S7)	Dec 12, 2014	ND (3.1)	9.1	9.1
Stingray (muscle)	Around 4km Offshore of Kumagawa (T-S8)	Dec 20, 2014	5.4	17	22.4
Blue crab (whole)	Around 4km Offshore of Kumagawa (T-S8)	Dec 20, 2014	ND (3.2)	ND (4.1)	ND

* When analyzed result is less than a detection limit level of the radioactivity concentration, showed as "ND"; as of individual Nuclide, detection limit level is showed in parenthesis.

* The sum of Cs-134 and Cs-137 radioactivity concentration as a standard value (since April 1, 2012) is 100 Bq/kg.

* Analyzed by Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shelfish (Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <5/11>
(excluding inside the port)

(Data summarized on January 20)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Snailfish (muscle)	Around 4km Offshore of Kumagawa (T-S8)	Dec 20, 2014	ND (3.8)	ND (4.1)	ND
Common skate (muscle)	Around 4km Offshore of Kumagawa (T-S8)	Dec 20, 2014	6.8	22	28.8
Microstomus achne (muscle)	Around 4km Offshore of Kumagawa (T-S8)	Dec 20, 2014	11	34	45
Ovalipes punctatus (whole)	Around 4km Offshore of Kumagawa (T-S8)	Dec 20, 2014	ND (4.9)	ND (3.8)	ND
Flatfish (muscle)	Around 4km Offshore of Kumagawa (T-S8)	Dec 20, 2014	ND (3.4)	5.6	5.6
Sea robin (muscle)	Around 4km Offshore of Kumagawa (T-S8)	Dec 20, 2014	ND (3.3)	ND (3.2)	ND
Marbled sole (muscle)	Around 4km Offshore of Kumagawa (T-S8)	Dec 20, 2014	5.7	22	27.7
Greenling (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Dec 16, 2014	ND (4.1)	5.5	5.5
Stone flounder (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Dec 16, 2014	ND (3.6)	6.0	6.0
Lepidotrigla microptera (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Dec 16, 2014	ND (3.6)	ND (3.9)	ND

* When analyzed result is less than a detection limit level of the radioactivity concentration, showed as "ND"; as of individual Nuclide, detection limit level is showed in parenthesis.

* The sum of Cs-134 and Cs-137 radioactivity concentration as a standard value (since April 1, 2012) is 100 Bq/kg.

* Analyzed by Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shelfish (Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <6/11>
(excluding inside the port)

(Data summarized on January 20)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Common skate (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Dec 16, 2014	ND (3.8)	11	11
Sea bass (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Dec 16, 2014	ND (4.4)	14	14
Crimson sea bream (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Dec 16, 2014	ND (3.8)	ND (4.1)	ND
Flatfish (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Dec 16, 2014	ND (3.9)	ND (4.1)	ND
Common Japanese conger (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Dec 16, 2014	ND (3.8)	ND (4.2)	ND
Littlemouth flounder (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Dec 16, 2014	ND (3.5)	4.1	4.1
Marbled sole (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Dec 16, 2014	ND (3.6)	4.8	4.8
Dory (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Dec 16, 2014	ND (3.4)	5.4	5.4
Roundnose flounder (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Dec 16, 2014	ND (3.2)	8.7	8.7
Ridged-eye flounder (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Dec 16, 2014	ND (3.1)	ND (3.8)	ND

* When analyzed result is less than a detection limit level of the radioactivity concentration, showed as "ND"; as of individual Nuclide, detection limit level is showed in parenthesis.

* The sum of Cs-134 and Cs-137 radioactivity concentration as a standard value (since April 1, 2012) is 100 Bq/kg.

* Analyzed by Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shelfish (Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <7/11>
(excluding inside the port)

(Data summarized on January 20)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Greenling (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Dec 16, 2014	ND (4.2)	ND (3.7)	ND
Striped jewfish (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Dec 16, 2014	ND (3.4)	ND (3.8)	ND
Blue crab (whole)	Around 18km Offshore of Ukedo River (T-B2)	Dec 16, 2014	ND (4.5)	ND (4.4)	ND
Lepidotrigla microptera (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Dec 16, 2014	ND (3.5)	3.4	3.4
Lophius litilon (whole)	Around 18km Offshore of Ukedo River (T-B2)	Dec 16, 2014	ND (3.6)	ND (3.3)	ND
Sea bass (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Dec 16, 2014	ND (3.9)	ND (3.7)	ND
Crimson sea bream (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Dec 16, 2014	ND (3.4)	4.1	4.1
Microstomus achne (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Dec 16, 2014	ND (3.6)	4.2	4.2
Flatfish (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Dec 16, 2014	ND (3.6)	4.4	4.4
Smooth dogfish (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Dec 16, 2014	ND (3.9)	7.6	7.6

* When analyzed result is less than a detection limit level of the radioactivity concentration, showed as "ND"; as of individual Nuclide, detection limit level is showed in parenthesis.

* The sum of Cs-134 and Cs-137 radioactivity concentration as a standard value (since April 1, 2012) is 100 Bq/kg.

* Analyzed by Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shelfish (Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <8/11>
(excluding inside the port)

(Data summarized on January 20)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Common Japanese conger (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Dec 16, 2014	ND (3.6)	4.9	4.9
Littlemouth flounder (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Dec 16, 2014	ND (4.4)	ND (3.7)	ND
Marbled sole (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Dec 16, 2014	ND (3.8)	4.8	4.8
Dory (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Dec 16, 2014	ND (3.9)	ND (3.9)	ND
Roundnose flounder (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Dec 16, 2014	ND (3.8)	ND (3.8)	ND
Ridged-eye flounder (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Dec 16, 2014	ND (4.3)	7.0	7.0
Stone flounder (muscle)	Around 10km Offshore of Fukushima Daiichi NPS (T-B3)	Dec 25, 2014	ND (4.1)	4.1	4.1
Lepidotrigla microptera (muscle)	Around 10km Offshore of Fukushima Daiichi NPS (T-B3)	Dec 25, 2014	ND (3.7)	ND (3.2)	ND
Common skete (muscle)	Around 10km Offshore of Fukushima Daiichi NPS (T-B3)	Dec 25, 2014	4.6	17	21.6
Sea bass (muscle)	Around 10km Offshore of Fukushima Daiichi NPS (T-B3)	Dec 25, 2014	ND (4.5)	ND (4.2)	ND

* When analyzed result is less than a detection limit level of the radioactivity concentration, showed as "ND"; as of individual Nuclide, detection limit level is showed in parenthesis.

* The sum of Cs-134 and Cs-137 radioactivity concentration as a standard value (since April 1, 2012) is 100 Bq/kg.

* Analyzed by Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shelfish (Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <9/11>
(excluding inside the port)

(Data summarized on January 20)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Crimson sea bream (muscle)	Around 10km Offshore of Fukushima Daiichi NPS (T-B3)	Dec 25, 2014	ND (4.1)	ND (3.5)	ND
Flatfish (muscle)	Around 10km Offshore of Fukushima Daiichi NPS (T-B3)	Dec 25, 2014	ND (3.4)	9.2	9.2
Sea robin (muscle)	Around 10km Offshore of Fukushima Daiichi NPS (T-B3)	Dec 25, 2014	ND (3.6)	3.8	3.8
Littlemouth flounder (muscle)	Around 10km Offshore of Fukushima Daiichi NPS (T-B3)	Dec 25, 2014	ND (3.6)	ND (3.6)	ND
Marbled sole (muscle)	Around 10km Offshore of Fukushima Daiichi NPS (T-B3)	Dec 25, 2014	ND (3.9)	ND (4.0)	ND
Roundnose flounder (muscle)	Around 10km Offshore of Fukushima Daiichi NPS (T-B3)	Dec 25, 2014	ND (4.1)	ND (4.2)	ND
Ridged-eye flounder (muscle)	Around 10km Offshore of Fukushima Daiichi NPS (T-B3)	Dec 25, 2014	ND (3.6)	4.9	4.9
Stone flounder (muscle)	Around 10km Offshore of Daini NPS (T-B4)	Dec 25, 2014	ND (3.8)	4.7	4.7
Lepidotrigla microptera (muscle)	Around 10km Offshore of Daini NPS (T-B4)	Dec 25, 2014	ND (3.7)	ND (4.1)	ND
Common skete (muscle)	Around 10km Offshore of Daini NPS (T-B4)	Dec 25, 2014	6.0	15	21

* When analyzed result is less than a detection limit level of the radioactivity concentration, showed as "ND"; as of individual Nuclide, detection limit level is showed in parenthesis.

* The sum of Cs-134 and Cs-137 radioactivity concentration as a standard value (since April 1, 2012) is 100 Bq/kg.

* Analyzed by Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shelfish (Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <10/11>
(excluding inside the port)

(Data summarized on January 20)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Sea bass (muscle)	Around 10km Offshore of Daini NPS (T-B4)	Dec 25, 2014	ND (3.1)	ND (3.8)	ND
Crimson sea bream (muscle)	Around 10km Offshore of Daini NPS (T-B4)	Dec 25, 2014	ND (3.3)	ND (3.8)	ND
Microstomus achne (muscle)	Around 10km Offshore of Daini NPS (T-B4)	Dec 25, 2014	5.2	25	30.2
Flatfish (muscle)	Around 10km Offshore of Daini NPS (T-B4)	Dec 25, 2014	ND (3.7)	5.2	5.2
Sea robin (muscle)	Around 10km Offshore of Daini NPS (T-B4)	Dec 25, 2014	ND (3.5)	ND (3.8)	ND
Smooth dogfish (muscle)	Around 10km Offshore of Daini NPS (T-B4)	Dec 25, 2014	ND (3.0)	ND (3.6)	ND
Common Japanese conger (muscle)	Around 10km Offshore of Daini NPS (T-B4)	Dec 25, 2014	ND (2.8)	5.1	5.1
Littlemouth flounder (muscle)	Around 10km Offshore of Daini NPS (T-B4)	Dec 25, 2014	ND (3.6)	5.4	5.4
Marbled sole (muscle)	Around 10km Offshore of Daini NPS (T-B4)	Dec 25, 2014	ND (3.5)	8.9	8.9
Roundnose flounder (muscle)	Around 10km Offshore of Daini NPS (T-B4)	Dec 25, 2014	ND (3.6)	ND (4.6)	ND

* When analyzed result is less than a detection limit level of the radioactivity concentration, showed as "ND"; as of individual Nuclide, detection limit level is showed in parenthesis.

* The sum of Cs-134 and Cs-137 radioactivity concentration as a standard value (since April 1, 2012) is 100 Bq/kg.

* Analyzed by Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shelfish (Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <11/11>
(excluding inside the port)

(Data summarized on January 20)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Ridged-eye flounder (muscle)	Around 10km Offshore of Daini NPS (T-B4)	Dec 25, 2014	ND (3.6)	ND (4.2)	ND

* When analyzed result is less than a detection limit level of the radioactivity concentration, showed as "ND"; as of individual Nuclide, detection limit level is showed in parenthesis.

* The sum of Cs-134 and Cs-137 radioactivity concentration as a standard value (since April 1, 2012) is 100 Bq/kg.

* Analyzed by Tokyo Power Technology Ltd.