

Nuclide Analysis Results of Fish and Shellfish (Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <1/10>
(excluding the Port Area)

(Data summarized on February 17)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Schlegel's black rockfish (muscle)	Around 1km Offshore of Ota River (T-S1)	Jan 22, 2016	ND(3.8)	ND(3.3)	ND
Common skate (muscle)	Around 1km Offshore of Ota River (T-S1)	Jan 22, 2016	ND(3.9)	8.7	8.7
Ovalipes punctatus (whole)	Around 1km Offshore of Ota River (T-S1)	Jan 22, 2016	ND(3.7)	ND(3.6)	ND
Flatfish (muscle)	Around 1km Offshore of Ota River (T-S1)	Jan 22, 2016	ND(4.1)	ND(4.2)	ND
Marbled sole (muscle)	Around 1km Offshore of Ota River (T-S1)	Jan 22, 2016	ND(3.7)	12	12
Pacific cod (muscle)	Around 1km Offshore of Ota River (T-S1)	Jan 22, 2016	ND(4.0)	ND(3.3)	ND
Blue crab (whole)	Around 3km Offshore of Odaka Ward (T-S2)	Jan 22, 2016	ND(4.1)	ND(3.9)	ND
Snailfish (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	Jan 22, 2016	ND(3.9)	ND(3.7)	ND
Schlegel's black rockfish (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	Jan 22, 2016	ND(4.0)	9.7	9.7
Common skate (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	Jan 22, 2016	ND(4.0)	15	15

* When an analysis result is less than a detection limit of radioactivity concentration, the value is indicated as "ND (not detected)."

A detection limit of individual nuclide is provided in parenthesis.

* The sum of Cs-134 and Cs-137 radioactivity concentration as a standard value (since April 1, 2012) is 100 Bq/kg.

* The samples are analyzed by Tokyo Power Technology, Ltd.

Nuclide Analysis Results of Fish and Shellfish (Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <2/10>
(excluding the Port Area)

(Data summarized on February 17)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Ovalipes punctatus (whole)	Around 3km Offshore of Odaka Ward (T-S2)	Jan 22, 2016	ND(3.7)	3.7	3.7
Flatfish (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	Jan 22, 2016	ND(3.3)	ND(3.8)	ND
Marbled sole (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	Jan 22, 2016	ND(3.7)	7.5	7.5
Pacific cod (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	Jan 22, 2016	ND(3.9)	ND(3.7)	ND
Snailfish (muscle)	Around 3km Offshore of Ukedo River (T-S3)	Jan 28, 2016	ND(3.3)	3.5	3.5
Schlegel's black rockfish (muscle)	Around 3km Offshore of Ukedo River (T-S3)	Jan 28, 2016	6.1	27	33.1
Common skate (muscle)	Around 3km Offshore of Ukedo River (T-S3)	Jan 28, 2016	ND(4.0)	13	13
Sea bass (muscle)	Around 3km Offshore of Ukedo River (T-S3)	Jan 28, 2016	ND(3.9)	ND(3.4)	ND
Ovalipes punctatus (whole)	Around 3km Offshore of Ukedo River (T-S3)	Jan 28, 2016	ND(3.3)	ND(3.3)	ND
Flatfish (muscle)	Around 3km Offshore of Ukedo River (T-S3)	Jan 28, 2016	ND(3.2)	6.0	6.0

* When an analysis result is less than a detection limit of radioactivity concentration, the value is indicated as "ND (not detected)."

A detection limit of individual nuclide is provided in parenthesis.

* The sum of Cs-134 and Cs-137 radioactivity concentration as a standard value (since April 1, 2012) is 100 Bq/kg.

* The samples are analyzed by Tokyo Power Technology, Ltd.

Nuclide Analysis Results of Fish and Shellfish (Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <3/10>
(excluding the Port Area)

(Data summarized on February 17)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Littlemouth flounder (muscle)	Around 3km Offshore of Ukedo River (T-S3)	Jan 28, 2016	ND(4.3)	ND(3.8)	ND
Marbled sole (muscle)	Around 3km Offshore of Ukedo River (T-S3)	Jan 28, 2016	ND(4.0)	9.4	9.4
Pacific cod (muscle)	Around 3km Offshore of Ukedo River (T-S3)	Jan 28, 2016	ND(3.7)	ND(4.1)	ND
Sebastes vulpes (muscle)	Around 3km Offshore of Fukushima Daiichi NPS (T-S4)	Jan 28, 2016	6.2	25	31.2
Common skate (muscle)	Around 3km Offshore of Fukushima Daiichi NPS (T-S4)	Jan 28, 2016	ND(4.1)	19	19
Microstomus achne (muscle)	Around 3km Offshore of Fukushima Daiichi NPS (T-S4)	Jan 28, 2016	5.4	27	32.4
Marbled sole (muscle)	Around 3km Offshore of Fukushima Daiichi NPS (T-S4)	Jan 28, 2016	ND(4.0)	9.8	9.8
Pacific cod (muscle)	Around 3km Offshore of Fukushima Daiichi NPS (T-S4)	Jan 28, 2016	ND(4.3)	ND(3.2)	ND
Roundnose flounder (muscle)	Around 3km Offshore of Fukushima Daiichi NPS (T-S4)	Jan 28, 2016	ND(3.8)	ND(3.5)	ND
Schlegel's black rockfish (muscle)	Around 2km Offshore of Kido River (T-S5)	Jan 27, 2016	5.4	29	34.4

* When an analysis result is less than a detection limit of radioactivity concentration, the value is indicated as "ND (not detected)."

A detection limit of individual nuclide is provided in parenthesis.

* The sum of Cs-134 and Cs-137 radioactivity concentration as a standard value (since April 1, 2012) is 100 Bq/kg.

* The samples are analyzed by Tokyo Power Technology, Ltd.

Nuclide Analysis Results of Fish and Shellfish (Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <4/10>
(excluding the Port Area)

(Data summarized on February 17)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Sea raven (muscle)	Around 2km Offshore of Kido River (T-S5)	Jan 27, 2016	ND(3.8)	ND(3.6)	ND
Common skete (muscle)	Around 2km Offshore of Kido River (T-S5)	Jan 27, 2016	ND(4.4)	21	21
Microstomus achne (muscle)	Around 2km Offshore of Kido River (T-S5)	Jan 27, 2016	ND(3.9)	7.9	7.9
Sea raven (muscle)	Offshore 2km vicinity of Fukushima Daini NPS (T-S7)	Jan 27, 2016	ND(3.7)	7.0	7.0
Common skete (muscle)	Offshore 2km vicinity of Fukushima Daini NPS (T-S7)	Jan 27, 2016	ND(3.2)	15	15
Microstomus achne (muscle)	Offshore 2km vicinity of Fukushima Daini NPS (T-S7)	Jan 27, 2016	5.8	27	32.8
Common skete (muscle)	Around 4km Offshore of Kumagawa (T-S8)	Jan 12, 2016	ND(3.2)	8.8	8.8
Microstomus achne (muscle)	Around 4km Offshore of Kumagawa (T-S8)	Jan 12, 2016	ND(3.9)	5.7	5.7
Flatfish (muscle)	Around 4km Offshore of Kumagawa (T-S8)	Jan 12, 2016	ND(3.6)	3.7	3.7
Sea robin (muscle)	Around 4km Offshore of Kumagawa (T-S8)	Jan 12, 2016	ND(3.9)	ND(3.7)	ND

* When an analysis result is less than a detection limit of radioactivity concentration, the value is indicated as "ND (not detected)."

A detection limit of individual nuclide is provided in parenthesis.

* The sum of Cs-134 and Cs-137 radioactivity concentration as a standard value (since April 1, 2012) is 100 Bq/kg.

* The samples are analyzed by Tokyo Power Technology, Ltd.

Nuclide Analysis Results of Fish and Shellfish (Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <5/10>
(excluding the Port Area)

(Data summarized on February 17)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Littlemouth flounder (muscle)	Around 4km Offshore of Kumagawa (T-S8)	Jan 12, 2016	ND(4.3)	4.9	4.9
Pacific cod (muscle)	Around 4km Offshore of Kumagawa (T-S8)	Jan 12, 2016	ND(3.6)	ND(4.1)	ND
Roundnose flounder (muscle)	Around 4km Offshore of Kumagawa (T-S8)	Jan 12, 2016	ND(3.3)	ND(4.2)	ND
Stone flounder (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Jan 14, 2016	ND(3.3)	7.8	7.8
Black scraper (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Jan 14, 2016	ND(4.0)	ND(3.3)	ND
Lepidotrigla microptera (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Jan 14, 2016	ND(3.8)	ND(4.0)	ND
Common skete (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Jan 14, 2016	ND(3.6)	11	11
Sea bass (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Jan 14, 2016	ND(4.2)	4.5	4.5
Microstomus achne (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Jan 14, 2016	ND(3.6)	ND(3.4)	ND
Flatfish (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Jan 14, 2016	ND(3.5)	3.9	3.9

* When an analysis result is less than a detection limit of radioactivity concentration, the value is indicated as "ND (not detected)."

A detection limit of individual nuclide is provided in parenthesis.

* The sum of Cs-134 and Cs-137 radioactivity concentration as a standard value (since April 1, 2012) is 100 Bq/kg.

* The samples are analyzed by Tokyo Power Technology, Ltd.

Nuclide Analysis Results of Fish and Shellfish (Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <6/10>
(excluding the Port Area)

(Data summarized on February 17)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Littlemouth flounder (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Jan 14, 2016	ND(4.2)	ND(4.2)	ND
Roundnose flounder (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Jan 14, 2016	ND(3.5)	ND(3.5)	ND
Greenling (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Jan 14, 2016	ND(3.8)	ND(4.4)	ND
Stone flounder (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Jan 14, 2016	ND(3.9)	ND(4.5)	ND
Lepidotrigla microptera (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Jan 14, 2016	ND(3.6)	ND(3.6)	ND
Lophius litilon(whole)	Around 18km Offshore of Ukedo River (T-B2)	Jan 14, 2016	ND(4.0)	ND(3.1)	ND
Common skete (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Jan 14, 2016	3.8	17	20.8
Microstomus achne (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Jan 14, 2016	ND(3.8)	ND(3.8)	ND
Flatfish (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Jan 14, 2016	ND(3.5)	5.3	5.3
Sea robin (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Jan 14, 2016	ND(3.3)	ND(3.7)	ND

* When an analysis result is less than a detection limit of radioactivity concentration, the value is indicated as "ND (not detected)."

A detection limit of individual nuclide is provided in parenthesis.

* The sum of Cs-134 and Cs-137 radioactivity concentration as a standard value (since April 1, 2012) is 100 Bq/kg.

* The samples are analyzed by Tokyo Power Technology, Ltd.

Nuclide Analysis Results of Fish and Shellfish (Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <7/10>
(excluding the Port Area)

(Data summarized on February 17)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Smooth dogfish (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Jan 14, 2016	ND(3.7)	ND(3.5)	ND
Littlemouth flounder (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Jan 14, 2016	ND(4.6)	4.0	4.0
Marbled sole (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Jan 14, 2016	ND(3.5)	3.9	3.9
Pacific cod (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Jan 14, 2016	ND(4.1)	ND(3.4)	ND
Roundnose flounder (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Jan 14, 2016	ND(3.9)	ND(4.2)	ND
Ridged-eye flounder (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Jan 14, 2016	ND(3.8)	3.3	3.3
Slippery sole (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Jan 14, 2016	ND(3.3)	ND(4.1)	ND
Greenling (muscle)	Around 10km Offshore of Fukushima Daiichi NPS (T-B3)	Jan 15, 2016	ND(4.4)	3.7	3.7
Stone flounder (muscle)	Around 10km Offshore of Fukushima Daiichi NPS (T-B3)	Jan 15, 2016	ND(4.4)	ND(2.9)	ND
Lepidotrigla microptera (muscle)	Around 10km Offshore of Fukushima Daiichi NPS (T-B3)	Jan 15, 2016	ND(3.8)	ND(3.8)	ND

* When an analysis result is less than a detection limit of radioactivity concentration, the value is indicated as "ND (not detected)."

A detection limit of individual nuclide is provided in parenthesis.

* The sum of Cs-134 and Cs-137 radioactivity concentration as a standard value (since April 1, 2012) is 100 Bq/kg.

* The samples are analyzed by Tokyo Power Technology, Ltd.

Nuclide Analysis Results of Fish and Shellfish (Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <8/10>
(excluding the Port Area)

(Data summarized on February 17)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Common skete (muscle)	Around 10km Offshore of Fukushima Daiichi NPS (T-B3)	Jan 15, 2016	ND(4.6)	12	12
Microstomus achne (muscle)	Around 10km Offshore of Fukushima Daiichi NPS (T-B3)	Jan 15, 2016	ND(4.0)	9.4	9.4
Flatfish (muscle)	Around 10km Offshore of Fukushima Daiichi NPS (T-B3)	Jan 15, 2016	ND(3.5)	ND(3.6)	ND
Sea robin (muscle)	Around 10km Offshore of Fukushima Daiichi NPS (T-B3)	Jan 15, 2016	ND(3.4)	ND(4.1)	ND
Smooth dogfish (muscle)	Around 10km Offshore of Fukushima Daiichi NPS (T-B3)	Jan 15, 2016	ND(3.9)	ND(3.6)	ND
Common Japanese conger (muscle)	Around 10km Offshore of Fukushima Daiichi NPS (T-B3)	Jan 15, 2016	ND(4.3)	ND(3.6)	ND
Littlemouth flounder(muscle)	Around 10km Offshore of Fukushima Daiichi NPS (T-B3)	Jan 15, 2016	ND(3.2)	ND(3.5)	ND
Marbled sole (muscle)	Around 10km Offshore of Fukushima Daiichi NPS (T-B3)	Jan 15, 2016	ND(3.7)	3.6	3.6
Roundnose flounder (muscle)	Around 10km Offshore of Fukushima Daiichi NPS (T-B3)	Jan 15, 2016	ND(3.6)	ND(4.0)	ND
Ridged-eye flounder (muscle)	Around 10km Offshore of Fukushima Daiichi NPS (T-B3)	Jan 15, 2016	ND(3.0)	ND(3.7)	ND

* When an analysis result is less than a detection limit of radioactivity concentration, the value is indicated as "ND (not detected)."

A detection limit of individual nuclide is provided in parenthesis.

* The sum of Cs-134 and Cs-137 radioactivity concentration as a standard value (since April 1, 2012) is 100 Bq/kg.

* The samples are analyzed by Tokyo Power Technology, Ltd.

Nuclide Analysis Results of Fish and Shellfish (Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <9/10>
(excluding the Port Area)

(Data summarized on February 17)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Stone flounder (muscle)	Around 10km Offshore of Fukushima Daini NPS (T-B4)	Jan 15, 2016	ND(3.8)	ND(3.6)	ND
Lepidotrigla microptera (muscle)	Around 10km Offshore of Fukushima Daini NPS (T-B4)	Jan 15, 2016	ND(3.5)	ND(4.4)	ND
Common skete(muscle)	Around 10km Offshore of Fukushima Daini NPS (T-B4)	Jan 15, 2016	ND(4.2)	14	14
Sea bass (muscle)	Around 10km Offshore of Fukushima Daini NPS (T-B4)	Jan 15, 2016	ND(4.0)	ND(3.6)	ND
Microstomus achne (muscle)	Around 10km Offshore of Fukushima Daini NPS (T-B4)	Jan 15, 2016	ND(3.1)	6.2	6.2
Flatfish (muscle)	Around 10km Offshore of Fukushima Daini NPS (T-B4)	Jan 15, 2016	ND(3.2)	ND(5.0)	ND
Sea robin (muscle)	Around 10km Offshore of Fukushima Daini NPS (T-B4)	Jan 15, 2016	ND(3.5)	ND(4.1)	ND
Smooth dogfish (muscle)	Around 10km Offshore of Fukushima Daini NPS (T-B4)	Jan 15, 2016	ND(3.5)	ND(3.5)	ND
Littlemouth flounder (muscle)	Around 10km Offshore of Fukushima Daini NPS (T-B4)	Jan 15, 2016	ND(2.9)	ND(3.6)	ND
Marbled sole (muscle)	Around 10km Offshore of Fukushima Daini NPS (T-B4)	Jan 15, 2016	ND(3.3)	8.0	8.0

* When an analysis result is less than a detection limit of radioactivity concentration, the value is indicated as "ND (not detected)."

A detection limit of individual nuclide is provided in parenthesis.

* The sum of Cs-134 and Cs-137 radioactivity concentration as a standard value (since April 1, 2012) is 100 Bq/kg.

* The samples are analyzed by Tokyo Power Technology, Ltd.

Nuclide Analysis Results of Fish and Shellfish (Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <10/10>
(excluding the Port Area)

(Data summarized on February 17)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Roundnose flounder (muscle)	Around 10km Offshore of Fukushima Daini NPS (T-B4)	Jan 15, 2016	ND(3.7)	ND(3.3)	ND
Ridged-eye flounder (muscle)	Around 10km Offshore of Fukushima Daini NPS (T-B4)	Jan 15, 2016	ND(3.1)	ND(3.2)	ND

* When an analysis result is less than a detection limit of radioactivity concentration, the value is indicated as "ND (not detected)."

A detection limit of individual nuclide is provided in parenthesis.

* The sum of Cs-134 and Cs-137 radioactivity concentration as a standard value (since April 1, 2012) is 100 Bq/kg.

* The samples are analyzed by Tokyo Power Technology, Ltd.