

## Nuclide Analysis Results of Fish <Port Area of Fukushima Daiichi Nuclear Power Station>

(Data summarized on January 19)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Schlegel's black rockfish (muscle)	Port area of Fukushima Daiichi NPS (Near the port entrance)	Dec. 3, 2016	30	180	210
Shad (muscle) No.1	Port area of Fukushima Daiichi NPS (Near the port entrance)	Dec. 3, 2016	ND(6.6)	13	13
Sebastes cheni (muscle)	Port area of Fukushima Daiichi NPS (Near the port entrance)	Dec. 3, 2016	1200	7100	8300
Striped mullet (muscle)	Port area of Fukushima Daiichi NPS (Near the port entrance)	Dec. 3, 2016	ND(4.3)	18	18
Greenling (muscle)	Port area of Fukushima Daiichi NPS (Near the port entrance)	Dec. 12, 2016	11	96	107
Acanthopagrus schlegeli (muscle)	Port area of Fukushima Daiichi NPS (Near the port entrance)	Dec. 12, 2016	14	76	90
Shad (muscle) No.2	Port area of Fukushima Daiichi NPS (Near the port entrance)	Dec. 12, 2016	ND(4.6)	8.3	8.3
Flatfish (muscle) No.1	Port area of Fukushima Daiichi NPS (Near the port entrance)	Dec. 12, 2016	140	780	920
Flatfish (muscle) No.2	Port area of Fukushima Daiichi NPS (Near the port entrance)	Dec. 17, 2016	ND(6.0)	33	33

\*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

\*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.