

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <1/12>
(excluding the port)

(Data summarized on January 19)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Greenling (muscle)	Around 1km Offshore of Ota River (T-S1)	Dec. 16, 2016	ND(4.2)	ND(4.6)	ND
Stingray (muscle)	Around 1km Offshore of Ota River (T-S1)	Dec. 16, 2016	ND(3.5)	ND(3.6)	ND
Blue crab (whole)	Around 1km Offshore of Ota River (T-S1)	Dec. 16, 2016	ND(4.1)	ND(3.3)	ND
Angel shark (muscle)	Around 1km Offshore of Ota River (T-S1)	Dec. 16, 2016	ND(4.3)	9.3	9.3
Schlegel's black rockfish (muscle)	Around 1km Offshore of Ota River (T-S1)	Dec. 16, 2016	3.8	22	25.8
Sea raven (muscle)	Around 1km Offshore of Ota River (T-S1)	Dec. 16, 2016	ND(3.3)	ND(4.1)	ND
Common skete (muscle)	Around 1km Offshore of Ota River (T-S1)	Dec. 16, 2016	ND(3.6)	ND(4.0)	ND
Microstomus achne (muscle)	Around 1km Offshore of Ota River (T-S1)	Dec. 16, 2016	ND(3.7)	ND(4.0)	ND
Flatfish (muscle)	Around 1km Offshore of Ota River (T-S1)	Dec. 16, 2016	ND(3.9)	ND(3.8)	ND
Smooth dogfish (muscle)	Around 1km Offshore of Ota River (T-S1)	Dec. 16, 2016	ND(4.0)	4.8	4.8

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <2/12>
(excluding the port)

(Data summarized on January 19)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Marbled sole (muscle)	Around 1km Offshore of Ota River (T-S1)	Dec. 16, 2016	ND(3.1)	13	13
Greenling (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	Dec. 16, 2016	ND(4.1)	ND(4.2)	ND
Stone flounder (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	Dec. 16, 2016	ND(3.4)	ND(3.6)	ND
Blue crab (whole)	Around 3km Offshore of Odaka Ward (T-S2)	Dec. 16, 2016	ND(4.0)	ND(4.0)	ND
Lepidotrigla microptera (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	Dec. 16, 2016	ND(3.4)	ND(4.3)	ND
Lophius litilon (whole)	Around 3km Offshore of Odaka Ward (T-S2)	Dec. 16, 2016	ND(3.7)	ND(3.0)	ND
Sea raven (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	Dec. 16, 2016	ND(3.5)	4.7	4.7
Common skete (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	Dec. 16, 2016	ND(4.1)	ND(3.9)	ND
Sea bass (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	Dec. 16, 2016	ND(4.2)	ND(3.1)	ND
Microstomus achne (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	Dec. 16, 2016	ND(4.3)	12	12

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <3/12>
(excluding the port)

(Data summarized on January 19)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Flatfish (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	Dec. 12, 2016	ND(4.2)	ND(3.2)	ND
Smooth dogfish (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	Dec. 12, 2016	ND(3.5)	ND(4.4)	ND
Marbled sole (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	Dec. 12, 2016	ND(3.5)	6.9	6.9
Roundnose flounder (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	Dec. 12, 2016	ND(3.2)	ND(4.0)	ND
Stingray (muscle)	Around 3km Offshore of Ukedo River (T-S3)	Dec. 12, 2016	ND(4.0)	ND(4.0)	ND
Stone flounder (muscle)	Around 3km Offshore of Ukedo River (T-S3)	Dec. 12, 2016	ND(3.1)	ND(4.1)	ND
Blue crab (whole)	Around 3km Offshore of Ukedo River (T-S3)	Dec. 12, 2016	ND(4.0)	ND(3.9)	ND
Angel shark (muscle)	Around 3km Offshore of Ukedo River (T-S3)	Dec. 12, 2016	18	120	138
Common skate (muscle)	Around 3km Offshore of Ukedo River (T-S3)	Dec. 12, 2016	ND(3.0)	15	15
Flatfish (muscle)	Around 3km Offshore of Ukedo River (T-S3)	Dec. 12, 2016	ND(3.7)	ND(4.8)	ND

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <4/12>
(excluding the port)

(Data summarized on January 19)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Marbled sole (muscle)	Around 3km Offshore of Ukedo River (T-S3)	Dec. 16, 2016	ND(3.7)	4.1	4.1
Flathead (Platycephalus sp.) (muscle)	Around 3km Offshore of Ukedo River (T-S3)	Dec. 16, 2016	ND(2.3)	5.7	5.7
Lophius litilon (whole)	Around 3km Offshore of Fukushima Daiichi (T-S4)	Dec. 16, 2016	ND(3.9)	ND(3.7)	ND
Common skete (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	Dec. 16, 2016	ND(4.2)	7.5	7.5
Drumfish (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	Dec. 16, 2016	ND(3.9)	ND(3.3)	ND
Microstomus achne (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	Dec. 16, 2016	ND(3.7)	11	11
Flatfish (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	Dec. 16, 2016	ND(4.0)	8.2	8.2
Marbled sole (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	Dec. 16, 2016	ND(4.1)	6.9	6.9
Pagrus major (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	Dec. 16, 2016	ND(4.3)	ND(3.7)	ND
Greenling (muscle)	Around 2km Offshore of Kido River (T-S5)	Dec. 12, 2016	ND(3.6)	6.3	6.3

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <5/12>
(excluding the port)

(Data summarized on January 19)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Common skete (muscle)	Around 2km Offshore of Kido River (T-S5)	Dec. 12, 2016	ND(3.3)	5.6	5.6
Drumfish (muscle)	Around 2km Offshore of Kido River (T-S5)	Dec. 12, 2016	ND(3.1)	ND(3.6)	ND
Microstomus achne (muscle)	Around 2km Offshore of Kido River (T-S5)	Dec. 12, 2016	ND(3.4)	ND(4.2)	ND
Flatfish (muscle)	Around 2km Offshore of Kido River (T-S5)	Dec. 12, 2016	ND(3.8)	5.1	5.1
Sea robin (muscle)	Around 2km Offshore of Kido River (T-S5)	Dec. 12, 2016	ND(3.8)	ND(4.2)	ND
Dasyatis matsubara (muscle)	Around 2km Offshore of Kido River (T-S5)	Dec. 12, 2016	ND(3.9)	ND(2.9)	ND
Greenling (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	Dec. 12, 2016	ND(3.1)	4.9	4.9
Stone flounder (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	Dec. 12, 2016	ND(4.4)	ND(3.7)	ND
Common skete (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	Dec. 12, 2016	ND(3.7)	10	10
Sebastes cheni (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	Dec. 12, 2016	5.1	34	39.1

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <6/12>
(excluding the port)

(Data summarized on January 19)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Banded dogfish (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	Dec. 12, 2016	ND(3.3)	ND(3.7)	ND
Drumfish (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	Dec. 12, 2016	ND(3.3)	ND(3.6)	ND
Microstomus achne (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	Dec. 12, 2016	ND(3.4)	18	18
Flatfish (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	Dec. 12, 2016	ND(3.4)	4.0	4.0
Sea robin (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	Dec. 12, 2016	ND(3.5)	4.5	4.5
Smooth dogfish (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	Dec. 12, 2016	ND(4.0)	3.7	3.7
Roundnose flounder (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	Dec. 12, 2016	ND(3.4)	3.6	3.6
Stingray (muscle)	Around 4km Offshore of Kumagawa (T-S8)	Dec. 9, 2016	ND(2.8)	ND(3.5)	ND
Blue crab (whole)	Around 4km Offshore of Kumagawa (T-S8)	Dec. 9, 2016	ND(3.5)	ND(4.5)	ND
Schlegel's black rockfish (muscle)	Around 4km Offshore of Kumagawa (T-S8)	Dec. 9, 2016	ND(3.6)	ND(4.0)	ND

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <7/12>
(excluding the port)

(Data summarized on January 19)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Flatfish (muscle)	Around 4km Offshore of Kumagawa (T-S8)	Dec. 9, 2016	ND(4.0)	5.1	5.1
Sea robin (muscle)	Around 4km Offshore of Kumagawa (T-S8)	Dec. 9, 2016	ND(3.3)	ND(3.3)	ND
Roundnose flounder (muscle)	Around 4km Offshore of Kumagawa (T-S8)	Dec. 9, 2016	ND(3.5)	ND(3.0)	ND
Greenling (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Dec. 8, 2016	ND(4.4)	ND(3.5)	ND
Stone flounder (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Dec. 8, 2016	ND(3.4)	ND(4.1)	ND
Striped jewfish (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Dec. 8, 2016	ND(3.3)	ND(3.1)	ND
Lepidotrigla microptera (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Dec. 8, 2016	ND(3.7)	ND(3.4)	ND
Common skete (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Dec. 8, 2016	ND(3.0)	4.3	4.3
Sea bass (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Dec. 8, 2016	ND(3.2)	ND(3.4)	ND
Flatfish (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Dec. 8, 2016	ND(3.4)	ND(3.8)	ND

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <8/12>
(excluding the port)

(Data summarized on January 19)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Littlemouth flounder (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Dec. 8, 2016	ND(3.1)	4.6	4.6
Marbled sole (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Dec. 8, 2016	ND(3.8)	ND(4.3)	ND
Pagrus major (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Dec. 8, 2016	ND(3.9)	ND(3.6)	ND
Dory (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Dec. 8, 2016	ND(3.5)	ND(3.8)	ND
Roundnose flounder (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Dec. 8, 2016	ND(3.3)	ND(3.1)	ND
Ridged-eye flounder (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Dec. 8, 2016	ND(3.4)	ND(3.7)	ND
Greenling (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Dec. 8, 2016	ND(3.9)	ND(3.4)	ND
Striped jewfish (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Dec. 8, 2016	ND(3.7)	ND(3.2)	ND
Lepidotrigla microptera (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Dec. 8, 2016	ND(3.9)	ND(3.9)	ND
Lophius litilon (whole)	Around 18km Offshore of Ukedo River (T-B2)	Dec. 8, 2016	ND(3.4)	ND(4.0)	ND

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <9/12>
(excluding the port)

(Data summarized on January 19)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Common skete (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Dec. 8, 2016	ND(3.2)	13	13
Pennahia argentata (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Dec. 8, 2016	ND(3.5)	ND(4.0)	ND
Sea bass (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Dec. 8, 2016	ND(3.1)	ND(3.4)	ND
Microstomus achne (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Dec. 8, 2016	ND(3.9)	ND(3.6)	ND
Flatfish (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Dec. 8, 2016	ND(3.5)	ND(3.8)	ND
Smooth dogfish (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Dec. 8, 2016	ND(3.5)	ND(3.7)	ND
Common Japanese conger (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Dec. 8, 2016	ND(3.3)	ND(3.4)	ND
Littlemouth flounder (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Dec. 8, 2016	ND(3.6)	ND(3.7)	ND
Marbled sole (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Dec. 8, 2016	ND(3.7)	ND(3.8)	ND
Dory (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Dec. 8, 2016	ND(3.3)	ND(3.2)	ND

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <10/12>
(excluding the port)

(Data summarized on January 19)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Roundnose flounder (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Dec. 8, 2016	ND(3.4)	ND(3.8)	ND
Ridged-eye flounder (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Dec. 8, 2016	ND(3.3)	ND(3.9)	ND
Stingray (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	Dec. 5, 2016	ND(3.2)	4.4	4.4
Stone flounder (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	Dec. 5, 2016	ND(3.0)	ND(3.1)	ND
Lepidotrigla microptera (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	Dec. 5, 2016	ND(3.5)	ND(3.3)	ND
Common skate (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	Dec. 5, 2016	ND(3.1)	12	12
Takifugu snyderi (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	Dec. 5, 2016	ND(3.5)	ND(3.7)	ND
Sea bass (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	Dec. 5, 2016	ND(4.0)	ND(3.4)	ND
Crimson sea bream (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	Dec. 5, 2016	ND(2.8)	ND(4.3)	ND
Flatfish (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	Dec. 5, 2016	ND(3.3)	ND(4.0)	ND

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <11/12>
(excluding the port)

(Data summarized on January 19)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Sea robin (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	Dec. 5, 2016	ND(3.6)	ND(3.9)	ND
Littlemouth flounder (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	Dec. 5, 2016	ND(3.7)	ND(3.9)	ND
Marbled sole (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	Dec. 5, 2016	ND(4.3)	ND(4.0)	ND
Pagrus major (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	Dec. 5, 2016	ND(2.7)	3.9	3.9
Roundnose flounder (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	Dec. 5, 2016	ND(3.5)	ND(4.0)	ND
Lepidotrigla microptera (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	Dec. 5, 2016	ND(3.7)	ND(3.9)	ND
Common skete (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	Dec. 5, 2016	ND(3.8)	7.3	7.3
Takifugu snyderi (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	Dec. 5, 2016	ND(4.0)	3.2	3.2
Sea bass (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	Dec. 5, 2016	ND(4.0)	ND(4.1)	ND
Banded dogfish (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	Dec. 5, 2016	ND(3.8)	ND(3.3)	ND

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <12/12>
(excluding the port)

(Data summarized on January 19)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Flatfish (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	Dec. 5, 2016	ND(3.4)	ND(3.0)	ND
Sea robin (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	Dec. 5, 2016	ND(3.6)	ND(3.7)	ND
Littlemouth flounder (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	Dec. 5, 2016	ND(3.9)	ND(3.8)	ND
Marbled sole (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	Dec. 5, 2016	ND(3.8)	3.9	3.9
Pagrus major (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	Dec. 5, 2016	ND(3.6)	ND(3.5)	ND
Roundnose flounder (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	Dec. 5, 2016	ND(4.0)	ND(3.8)	ND

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.