

**Nuclide Analysis Results of Fish and Shellfish**  
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <1/12>  
(excluding the port)

(Data summarized on April 14)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Schlegel's black rockfish (muscle)	Around 1km Offshore of Ota River (T-S1)	Mar. 10, 2017	ND(3.7)	ND(4.0)	ND
Common skete (muscle)	Around 1km Offshore of Ota River (T-S1)	Mar. 10, 2017	ND(4.4)	19	19
Microstomus achne (muscle)	Around 1km Offshore of Ota River (T-S1)	Mar. 10, 2017	ND(4.1)	ND(3.9)	ND
Flatfish (muscle)	Around 1km Offshore of Ota River (T-S1)	Mar. 10, 2017	ND(4.0)	ND(3.3)	ND
Marbled sole (muscle)	Around 1km Offshore of Ota River (T-S1)	Mar. 10, 2017	ND(3.7)	ND(3.7)	ND
Stone flounder (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	Mar. 10, 2017	ND(3.6)	ND(3.9)	ND
Sebastes vulpes (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	Mar. 10, 2017	ND(3.2)	ND(3.9)	ND
Common skete (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	Mar. 10, 2017	ND(3.5)	5.7	5.7
Microstomus achne (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	Mar. 10, 2017	ND(3.4)	5.1	5.1
Littlemouth flounder (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	Mar. 10, 2017	ND(3.2)	ND(4.1)	ND

\*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

\*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

\*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish  
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <2/12>  
(excluding the port)

(Data summarized on April 14)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Marbled sole (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	Mar. 10, 2017	ND(3.9)	8.6	8.6
Pacific cod (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	Mar. 10, 2017	ND(4.4)	ND(3.7)	ND
Stingray (muscle)	Around 3km Offshore of Ukedo River (T-S3)	Mar. 9, 2017	ND(3.5)	ND(3.8)	ND
Stone flounder (muscle)	Around 3km Offshore of Ukedo River (T-S3)	Mar. 9, 2017	ND(3.6)	ND(3.2)	ND
Blue crab (whole)	Around 3km Offshore of Ukedo River (T-S3)	Mar. 9, 2017	ND(3.2)	4.2	4.2
Lepidotrigla microptera (muscle)	Around 3km Offshore of Ukedo River (T-S3)	Mar. 9, 2017	ND(3.1)	ND(3.2)	ND
Schlegel's black rockfish (muscle)	Around 3km Offshore of Ukedo River (T-S3)	Mar. 9, 2017	ND(3.8)	6.7	6.7
Common skete (muscle)	Around 3km Offshore of Ukedo River (T-S3)	Mar. 9, 2017	ND(3.4)	7.0	7.0
Microstomus achne (muscle)	Around 3km Offshore of Ukedo River (T-S3)	Mar. 9, 2017	ND(3.7)	9.1	9.1
Ovalipes punctatus (whole)	Around 3km Offshore of Ukedo River (T-S3)	Mar. 9, 2017	ND(4.4)	ND(3.7)	ND

\*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

\*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

\*Analyzed by: Tokyo Power Technology Ltd.

**Nuclide Analysis Results of Fish and Shellfish**  
**(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <3/12>**  
**(excluding the port)**

(Data summarized on April 14)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Flatfish① (muscle)	Around 3km Offshore of Ukedo River (T-S3)	Mar. 9, 2017	ND(3.6)	ND(3.8)	ND
Flatfish② (muscle)	Around 3km Offshore of Ukedo River (T-S3)	Mar. 9, 2017	ND(3.6)	ND(3.9)	ND
Smooth dogfish (muscle)	Around 3km Offshore of Ukedo River (T-S3)	Mar. 9, 2017	ND(4.3)	4.3	4.3
Marbled sole (muscle)	Around 3km Offshore of Ukedo River (T-S3)	Mar. 9, 2017	ND(4.1)	7.8	7.8
Pacific cod (muscle)	Around 3km Offshore of Ukedo River (T-S3)	Mar. 9, 2017	ND(4.5)	ND(4.3)	ND
Roundnose flounder (muscle)	Around 3km Offshore of Ukedo River (T-S3)	Mar. 9, 2017	ND(3.5)	ND(3.3)	ND
Frog flounder (muscle)	Around 3km Offshore of Ukedo River (T-S3)	Mar. 9, 2017	ND(3.7)	3.9	3.9
Stone flounder (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	Mar. 9, 2017	ND(3.5)	ND(3.7)	ND
Sebastes vulpes (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	Mar. 9, 2017	ND(3.7)	ND(3.9)	ND
Sea raven (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	Mar. 9, 2017	ND(4.0)	17	17

\*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

\*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

\*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish  
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <4/12>  
(excluding the port)

(Data summarized on April 14)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Common skate (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	Mar. 9, 2017	ND(3.9)	10	10
Sebastes cheni (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	Mar. 9, 2017	ND(3.1)	9.2	9.2
Pointhead flounder (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	Mar. 9, 2017	ND(3.8)	ND(3.4)	ND
Microstomus achne (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	Mar. 9, 2017	ND(3.2)	8.8	8.8
Ovalipes punctatus (whole)	Around 3km Offshore of Fukushima Daiichi (T-S4)	Mar. 9, 2017	ND(3.7)	ND(3.9)	ND
Flatfish① (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	Mar. 9, 2017	ND(3.8)	ND(4.6)	ND
Flatfish② (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	Mar. 9, 2017	ND(3.7)	4.4	4.4
Littlemouth flounder (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	Mar. 9, 2017	ND(4.5)	ND(3.1)	ND
Marbled sole (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	Mar. 9, 2017	ND(3.6)	3.8	3.8
Littlemouth flounder (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	Mar. 9, 2017	ND(3.3)	ND(3.7)	ND

\*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

\*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

\*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish  
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <5/12>  
(excluding the port)

(Data summarized on April 14)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Pacific cod (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	Mar. 9, 2017	ND(3.2)	ND(3.8)	ND
Roundnose flounder (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	Mar. 9, 2017	ND(3.7)	ND(4.3)	ND
Sea raven (muscle)	Around 2km Offshore of Kido River (T-S5)	Mar. 14, 2017	ND(4.1)	ND(3.5)	ND
Common skete (muscle)	Around 2km Offshore of Kido River (T-S5)	Mar. 14, 2017	ND(3.7)	15	15
Microstomus achne (muscle)	Around 2km Offshore of Kido River (T-S5)	Mar. 14, 2017	ND(4.3)	6.4	6.4
Marbled sole (muscle)	Around 2km Offshore of Kido River (T-S5)	Mar. 14, 2017	ND(3.0)	8.9	8.9
Sebastes vulpes (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	Mar. 14, 2017	ND(4.3)	18	18
Sea raven (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	Mar. 14, 2017	ND(3.6)	ND(3.6)	ND
Common skete (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	Mar. 14, 2017	ND(4.1)	6.2	6.2
Microstomus achne (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	Mar. 14, 2017	ND(4.0)	9.7	9.7

\*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

\*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

\*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish  
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <6/12>  
(excluding the port)

(Data summarized on April 14)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Marbled sole (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	Mar. 14, 2017	4.7	24	28.7
Roundnose flounder (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	Mar. 14, 2017	ND(4.1)	ND(3.8)	ND
Stingray (muscle)	Around 4km Offshore of Kumagawa (T-S8)	Mar. 7, 2017	ND(3.8)	ND(3.4)	ND
Common skete (muscle)	Around 4km Offshore of Kumagawa (T-S8)	Mar. 7, 2017	ND(3.3)	7.3	7.3
Microstomus achne (muscle)	Around 4km Offshore of Kumagawa (T-S8)	Mar. 7, 2017	ND(4.1)	10	10
Flatfish (muscle)	Around 4km Offshore of Kumagawa (T-S8)	Mar. 7, 2017	ND(4.1)	ND(4.4)	ND
Littlemouth flounder (muscle)	Around 4km Offshore of Kumagawa (T-S8)	Mar. 7, 2017	ND(3.5)	4.4	4.4
Marbled sole (muscle)	Around 4km Offshore of Kumagawa (T-S8)	Mar. 7, 2017	ND(3.5)	ND(4.0)	ND
Roundnose flounder (muscle)	Around 4km Offshore of Kumagawa (T-S8)	Mar. 7, 2017	ND(4.3)	ND(3.7)	ND
Greenling (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Mar. 30, 2017	ND(3.4)	ND(4.4)	ND

\*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

\*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

\*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish  
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <7/12>  
(excluding the port)

(Data summarized on April 14)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Stone flounder (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Mar. 30, 2017	ND(3.7)	ND(4.0)	ND
Lepidotrigla microptera (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Mar. 30, 2017	ND(3.1)	ND(3.9)	ND
Common skete (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Mar. 30, 2017	ND(3.8)	ND(4.3)	ND
Pointhead flounder (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Mar. 30, 2017	ND(3.5)	ND(4.0)	ND
Microstomus achne (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Mar. 30, 2017	ND(3.7)	ND(3.9)	ND
Flatfish① (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Mar. 30, 2017	ND(3.9)	ND(3.6)	ND
Flatfish② (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Mar. 30, 2017	ND(3.4)	5.2	5.2
Common Japanese conger (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Mar. 30, 2017	ND(3.0)	ND(3.8)	ND
Littlemouth flounder (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Mar. 30, 2017	ND(3.3)	ND(4.6)	ND
Marbled sole (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Mar. 30, 2017	ND(3.3)	ND(3.4)	ND

\*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

\*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

\*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish  
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <8/12>  
(excluding the port)

(Data summarized on April 14)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Pacific cod (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Mar. 30, 2017	ND(4.0)	ND(3.9)	ND
Roundnose flounder (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Mar. 30, 2017	ND(3.9)	ND(3.7)	ND
Greenling (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Mar. 30, 2017	ND(3.9)	ND(3.2)	ND
Lepidotrigla microptera (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Mar. 30, 2017	ND(3.7)	ND(2.8)	ND
Lophius litilon (whole)	Around 18km Offshore of Ukedo River (T-B2)	Mar. 30, 2017	ND(3.5)	ND(3.6)	ND
Common skate (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Mar. 30, 2017	ND(3.3)	ND(4.3)	ND
Sea bass (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Mar. 30, 2017	ND(3.4)	ND(3.9)	ND
Microstomus achne (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Mar. 30, 2017	ND(3.3)	ND(3.1)	ND
Flatfish① (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Mar. 30, 2017	ND(3.4)	ND(3.6)	ND
Flatfish② (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Mar. 30, 2017	ND(3.6)	ND(3.1)	ND

\*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

\*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

\*Analyzed by: Tokyo Power Technology Ltd.

**Nuclide Analysis Results of Fish and Shellfish**  
**(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <9/12>**  
**(excluding the port)**

(Data summarized on April 14)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Common Japanese conger (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Mar. 30, 2017	ND(3.5)	ND(3.2)	ND
Littlemouth flounder (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Mar. 30, 2017	ND(3.5)	ND(3.7)	ND
Marbled sole (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Mar. 30, 2017	ND(3.5)	ND(3.7)	ND
Pacific cod (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Mar. 30, 2017	ND(3.5)	ND(3.8)	ND
Roundnose flounder (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Mar. 30, 2017	ND(3.5)	ND(4.1)	ND
Ridged-eye flounder (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Mar. 30, 2017	ND(3.2)	ND(3.0)	ND
willowy flounder (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Mar. 30, 2017	ND(4.0)	ND(3.4)	ND
Greenling (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	Mar. 18, 2017	ND(4.5)	6.6	6.6
Stone flounder (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	Mar. 18, 2017	ND(2.9)	ND(4.0)	ND
Lepidotrigla microptera (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	Mar. 18, 2017	ND(4.3)	ND(3.3)	ND

\*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

\*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

\*Analyzed by: Tokyo Power Technology Ltd.

**Nuclide Analysis Results of Fish and Shellfish**  
**(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <10/12>**  
**(excluding the port)**

(Data summarized on April 14)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Common skate (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	Mar. 18, 2017	ND(4.3)	ND(4.6)	ND
Sea bass (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	Mar. 18, 2017	ND(3.6)	ND(3.7)	ND
Microstomus achne (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	Mar. 18, 2017	ND(3.9)	6.5	6.5
Flatfish① (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	Mar. 18, 2017	ND(4.1)	ND(3.1)	ND
Flatfish② (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	Mar. 18, 2017	ND(3.5)	4.5	4.5
Sea robin (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	Mar. 18, 2017	ND(3.5)	ND(3.9)	ND
Smooth dogfish (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	Mar. 18, 2017	ND(3.9)	ND(4.4)	ND
Littlemouth flounder (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	Mar. 18, 2017	ND(3.7)	ND(3.3)	ND
Marbled sole (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	Mar. 18, 2017	ND(2.8)	ND(3.8)	ND
Pacific cod (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	Mar. 18, 2017	ND(4.0)	ND(3.8)	ND

\*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

\*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

\*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish  
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <11/12>  
(excluding the port)

(Data summarized on April 14)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Lepidotrigla microptera (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	Mar. 18, 2017	ND(3.9)	ND(3.5)	ND
Common skete (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	Mar. 18, 2017	ND(3.7)	6.4	6.4
Pointhead flounder (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	Mar. 18, 2017	ND(4.3)	ND(3.9)	ND
Microstomus achne (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	Mar. 18, 2017	ND(4.1)	10	10
Flatfish① (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	Mar. 18, 2017	ND(3.6)	ND(3.8)	ND
Flatfish② (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	Mar. 18, 2017	ND(4.2)	ND(3.9)	ND
Smooth dogfish (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	Mar. 18, 2017	ND(3.8)	ND(3.6)	ND
Littlemouth flounder (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	Mar. 18, 2017	ND(3.3)	ND(3.1)	ND
Marbled sole (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	Mar. 18, 2017	ND(3.6)	4.2	4.2
Pacific cod (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	Mar. 18, 2017	ND(3.4)	ND(3.7)	ND

\*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

\*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

\*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish  
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <12/12>  
(excluding the port)

(Data summarized on April 14)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Roundnose flounder (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	Mar. 18, 2017	ND (4. 0)	ND (3. 2)	ND
Ridged-eye flounder (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	Mar. 18, 2017	ND (3. 9)	ND (3. 3)	ND

\*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

\*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

\*Analyzed by: Tokyo Power Technology Ltd.