

Nuclide Analysis Results of Fish <Port Area of Fukushima Daiichi Nuclear Power Station>

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Greenling (muscle)	Port area of Fukushima Daiichi NPS(In front of mega-float)	January 11, 2018	13	140	153
Striped mullet (muscle) No.1	Port area of Fukushima Daiichi NPS(In front of mega-float)	January 11, 2018	ND(7.0)	22	22
Striped mullet (muscle) No.2	Port area of Fukushima Daiichi NPS(In front of mega-float)	January 11, 2018	ND(6.0)	42	42
Striped mullet (muscle) No.3	Port area of Fukushima Daiichi NPS(In front of mega-float)	January 11, 2018	ND(5.5)	24	24
Striped mullet (muscle) No.4	Port area of Fukushima Daiichi NPS(In front of mega-float)	January 11, 2018	33	250	283
Striped mullet (muscle) No.5	Port area of Fukushima Daiichi NPS(In front of mega-float)	January 11, 2018	10	64	74
Black rockfish (muscle)	Port area of Fukushima Daiichi NPS(In front of silt fence)	January 11, 2018	19	160	179

*When analyzed results are less than detection limits of radioactivity concentrations, the values are shown as "ND." Detection limits of individual nuclides are shown in page 10.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.